

BINARY DISRUPTION BY MASSIVE BLACK HOLES IN GLOBULAR CLUSTERS

ERIC PFAHL

Chandra Fellow; Department of Astronomy, University of Virginia, P.O. Box 3818, Charlottesville, VA 22903-0818; epfahl@virginia.edu

Accepted by ApJ Letters

ABSTRACT

A massive ($\sim 10^2$ – $10^4 M_\odot$) black hole at the center of a globular cluster will tidally disrupt binaries that pass sufficiently close. Such an encounter results in the capture of one binary component by the black hole and ejection of the other star. A $10^3 M_\odot$ black hole disrupts one binary every 1–10 Myr; captured stars orbit with initial periods of days to decades. Although $\gtrsim 100$ stars are captured in only 1 Gyr, the number with periods of < 10 yr at any one time is probably less than 10, regulated by stellar interactions and tidal disruption by the black hole. Binary disruption and stellar capture should be critically examined with detailed numerical simulations, in order to corroborate the analytic results given here and to weigh possible observational consequences.

Subject headings: black hole physics — globular clusters: general — stellar dynamics

1. INTRODUCTION

Black holes of mass $\sim 10^2$ – $10^4 M_\odot$ occupy a special niche in several subfields of theoretical astrophysics. Such objects might naturally form as remnants of the first stars in the Universe (Barkana & Loeb 2001; Madau & Rees 2001; Schneider et al. 2002), or be built through merging of stars (Portegies Zwart & McMillan 2002) or stellar-mass black holes (Miller & Hamilton 2002) in young star clusters. These black holes may play critical roles in the formation and growth of supermassive ($\gtrsim 10^6 M_\odot$) black holes in galactic nuclei (Ebisuzaki et al. 2001; Islam, Taylor, & Silk 2004).

Intermediate-mass black holes (IMBHs) have been invoked to resolve several recent observational quandaries. Some of the “ultraluminous” X-ray sources detected in other galaxies may be accreting IMBHs (e.g., Miller & Colbert 2004), although the majority are likely stellar-mass black holes (King et al. 2001; Rappaport, Podsiadlowski, & Pfahl 2004). There have been recent claims of possible IMBH detections in two globular clusters (M15 in the Milky Way and G1 in M31) based on *HST* data (Gebhardt, Rich, & Ho 2002; Gerssen et al. 2002). However, the data do not require IMBHs, as shown by Baumgardt et al. (2003a,b) and McNamara, Harrison, & Anderson (2003). Colpi, Possenti, & Gualandris (2002) cite evidence for a single or binary IMBH in the cluster NGC 6752 based on the anomalous positions and accelerations of cluster radio pulsars. Finally, a single or binary IMBH could account for the net rotation observed in the center of M15 (Gebhardt et al. 2000; Gerssen et al. 2002; Miller & Colbert 2004).

Because of the broad scientific scope sampled by IMBHs, it is crucial to investigate what unique dynamical signatures they imprint on their environments. The high stellar densities and proximities of Galactic globular clusters make them prime locales to search for IMBHs, as emphasized by much of the recent observational work; for a review of early theoretical and observational studies, see Shapiro (1985). Here we introduce a process that deposits stars directly into compact orbits (e.g., $\lesssim 100$ AU) around a massive cluster black hole.

Binary stars are abundant in globular clusters (e.g., Hut et al. 1992). A binary that passes sufficiently near a central black hole is disrupted by tidal forces, resulting in the capture of one star and ejection of the other (see § 2). This type of interaction was studied in the context of Sgr A* by Hills (1988) and Yu & Tremaine (2003) as a mechanism for accel-

erating stars to very high speeds, and by Gould & Quillen (2003) to account for the young stars near Sgr A* (e.g., Eisenhauer et al. 2005). The disruption rate in globular clusters is estimated in § 3, and the fate of captured stars is discussed in § 4. Future directions for studying this mechanism in more detail are suggested in § 5, as are potentially observable consequences.

2. BINARY DISRUPTION

Consider a binary system of mass $M_b = M_1 + M_2$ with semi-major axis a_b on a parabolic approach to a black hole of mass $M_\bullet \gg M_b$. Let r be the distance between the binary barycenter and the black hole, and s be the *minimum* distance. The differential acceleration across the binary ($\sim GM_\bullet a_b / r^3$) exceeds the binary self-gravity ($\sim GM_b / a_b^2$) within the tidal radius,

$$r_t \equiv a_b (M_\bullet / M_b)^{1/3} = 10 a_b Q_3^{1/3}, \quad (1)$$

where $M_\bullet / M_b = 10^3 Q_3$. For a disrupted binary, the tidal perturbation is essentially impulsive, and the additional relative velocity imparted to the binary components has a magnitude of $\sim v_b = (GM_b / a_b)^{1/2}$. Immediately following the disruption, the two stars are mutually unbound and move relative to the black hole with the original systemic velocity of magnitude $u_t \simeq (2GM_\bullet / r_t)^{1/2}$, plus contributions from the prior orbital motion and tidal perturbation. Random orbital orientations and phases at the time of disruption lead to a range of orbital parameters and asymptotic speeds for stars that are captured or ejected; below are estimates for the typical values.

If star 1 is captured, we can estimate the expectation value of its specific energy, E_1 , relative to the black hole by assuming that its resultant velocity after the tidal perturbation is directed opposite to the binary systemic velocity and has a magnitude of $(M_2 / M_b) v_b$. To leading order, we obtain

$$E_1 \simeq \frac{1}{2} \left(u_t - \frac{M_2}{M_b} v_b \right)^2 - \frac{GM_\bullet}{r_t} \simeq -\frac{M_2}{M_b} u_t v_b. \quad (2)$$

The corresponding semimajor axis and orbital period are

$$a_1 = -GM_\bullet / 2E_1 \simeq 35 a_b (M_b / M_2) Q_3^{2/3}, \quad (3)$$

and

$$P_1 \simeq 7 P_b (M_b/M_2)^{3/2} Q_3^{1/2}, \quad (4)$$

where P_b is the binary period. The pericenter separation of star 1, $a_1(1-e_1)$, must be comparable to s , so that

$$e_1 \simeq 1 - \frac{s}{a_1} \simeq 1 - 0.3 \left(\frac{s}{r_t} \right) \left(\frac{M_2}{M_b} \right) Q_3^{-1/3}, \quad (5)$$

which is $\gtrsim 0.7$ for $Q_3 > 1$. Star 2 has a typical specific energy of $E_2 \simeq (M_1/M_b) v_t v_b$, and thus its speed at infinity is

$$v_2 \simeq 4 v_b (M_1/M_b)^{1/2} Q_3^{1/6}, \quad (6)$$

which may be $\gtrsim 100 \text{ km s}^{-1}$ for typical binary parameters. A star ejected by this process can easily escape its host cluster.

3. TOTAL DISRUPTION RATE

We now employ “loss-cone” theory to estimate the rate of binary disruptions. The approach taken here is not very rigorous, but captures the essential physics. More precise accounts of loss-cone dynamics can be found in, e.g., Frank & Rees (1976), Lightman & Shapiro (1977), Cohn & Kulsrud (1978), Syer & Ulmer (1999), and Magorrian & Tremaine (1999).

A binary with semimajor axis a_b at radius $r \gg r_t$ is tidally disrupted if its specific angular momentum about the black hole is less than $J_t = (2GM_\bullet r_t)^{1/2}$. If the local one-dimensional stellar velocity dispersion, $\sigma(r)$, is adopted as a typical binary systemic speed, the disruption condition defines a double-sided loss cone in velocity space with semiangle $\theta_t = J_t/r\sigma$. Perturbations by distant stars cause the binary barycenter to undergo a random walk in phase space. In a dynamical time, $\tau_d \sim r/\sigma$, the velocity is deflected by an angle $\theta_d \sim (\tau_d/\tau_r)^{1/2}$, measured at radius r . The local two-body relaxation timescale is given by (e.g., Spitzer & Hart 1971)

$$\tau_r(r) = \frac{\sigma^3(r)}{\Lambda G^2 m^2 n(r)} \simeq 50 \frac{(\sigma/10 \text{ km s}^{-1})^3}{\Lambda_{10} (n/10^5 \text{ pc}^{-3})} \text{ Myr}, \quad (7)$$

where $n(r)$ is the stellar number density, $\Lambda = \Lambda_{10} 10$ is proportional to the Coulomb logarithm, and we have assumed $m = 1 M_\odot$ for the mean stellar and binary mass, which is fixed in the subsequent calculations.

The ratio θ_d/θ_t increases monotonically with r for profiles $n(r)$ and $\sigma(r)$ suitable to globular clusters. Two loss-cone regimes are loosely demarcated by $\theta_d/\theta_t = 1$, defining a critical radius r_c . For $r > r_c$, binaries can scatter into and out of the loss cone in a dynamical time, and the loss cone is full statistically. If $r < r_c$, binaries within the loss cone are disrupted in a dynamical time, and the disruption rate is limited by the diffusion of binaries across the loss-cone boundary.

We assume that the cluster center consists of a power-law density cusp around the black hole, embedded in an isothermal cluster core of radius r_0 . The cusp is inside the radius of influence of the central black hole:

$$r_i = GM_\bullet/\sigma_i^2 \simeq 0.04 M_3 \sigma_{10}^{-2} \text{ pc}, \quad (8)$$

where $M_\bullet = 10^3 M_3 M_\odot$, and $\sigma_i = 10 \sigma_{10} \text{ km s}^{-1}$ is the velocity dispersion at r_i . Each zone has different power-law profiles,

$n(r) = n_i (r/r_i)^{-\alpha}$ and $\sigma^2(r) = \sigma_i^2 (r/r_i)^{-\beta}$, where $\alpha = \beta = 0$ in the core, and $\alpha \simeq 1.5-2$ (e.g., Bahcall & Wolf 1977; Young 1977) and $\beta = 1$ in the cusp. The critical radius is

$$r_c/r_i = (a_b/a_c)^{1/(3+\beta-\alpha)}, \quad (9)$$

where, given $n_i = 10^5 n_5 \text{ pc}^{-3}$, we have

$$a_c \simeq 0.1 n_5 M_3^{5/3} \sigma_{10}^{-8} \Lambda_{10} \text{ AU}. \quad (10)$$

Cluster binaries interact frequently with single stars and other binaries. These encounters shape the binary population as the cluster and its stars evolve, a complex problem that can only be fully unraveled with detailed cluster simulations (e.g., Fregeau et al. 2003; Giersz & Spurzem 2003). Here we assume for simplicity a differential distribution $p(a_b) \propto a_b^{-1}$ over the range $a_b = 0.01-1 \text{ AU}$, and a fixed binary number fraction f_b . As the cluster evolves, the shape and limits of $p(a_b)$ vary, and f_b in the core may decrease from $\gtrsim 0.5$ initially to current values of $\lesssim 0.1$ (Ivanova et al. 2005).

When the loss cone is full, a fraction $\simeq \theta_t^2$ of the binaries with semimajor axis a_b at radius r are disrupted in a dynamical time. The differential rate is

$$\frac{d\mathcal{R}_{\text{full}}}{da_b} \simeq 4\pi f_b GM_\bullet \left(\frac{M_\bullet}{m} \right)^{1/3} p(a_b) a_b \int_{r_c}^{r_0} \frac{dr}{r} \frac{n(r)}{\sigma(r)}. \quad (11)$$

The contribution from $r > r_i$ dominates, and we obtain

$$\mathcal{R}_{\text{full}} \sim 10^{-5} f_b n_5 \langle a_b(\text{AU}) \rangle M_3^{4/3} \sigma_{10}^{-1} \ln \left(\frac{r_0}{r_i} \right) \text{ yr}^{-1}, \quad (12)$$

where $\langle a_b(\text{AU}) \rangle$ is the mean semimajor axis in AU.

In the diffusive regime, the fraction of binaries disrupted in a time $\tau_r(r)$ is $\sim 1/\ln(2/\theta_t)$ (e.g., Frank & Rees 1976). The differential diffusive disruption rate is then

$$\frac{d\mathcal{R}_{\text{diff}}}{da_b} \simeq 4\pi f_b G^2 m^2 p(a_b) \int_{r_c}^{r_0} \frac{dr}{r} \frac{r^3 n^2(r)}{\sigma^3(r)}, \quad (13)$$

where the weakly varying factor $\Lambda/\ln(2/\theta_t) \sim 1$ has been dropped. The lower limit in the above radial integral is relatively unimportant if it is $\ll r_c$. Numerically, we find

$$\mathcal{R}_{\text{diff}} \sim 10^{-6} f_b n_5^2 M_3^3 \sigma_{10}^{-9} \text{ yr}^{-1}. \quad (14)$$

A binary fraction of $f_b \sim 0.1$ and mean semimajor axis of $\langle a_b \rangle \sim 0.1 \text{ AU}$ give disruption rates of $\sim 10^{-7} \text{ yr}^{-1}$ for both the full and diffusive regimes when $M_3 = n_5 = \sigma_{10} = 1$. The total disruption rate is then $\sim 10^{-7}-10^{-6} \text{ yr}^{-1}$ for the same parameter values. For $a_b = 0.01-1 \text{ AU}$ and $M_3 \simeq 1$, captured stars have $a_1 \sim 1-100 \text{ AU}$ when $M_1 \simeq M_2$, and periods of $\sim 10 \text{ d}$ to $\sim 30 \text{ yr}$. From eq. (3), $p(a_1)$ should have a similar shape as $p(a_b)$. Our choice of $p(a_b) \propto a_b^{-1}$ implies that the distributions of $\log a_1$, $\log P_1$, and $\log v_2$ should be roughly flat.

As an illustration, we computed 10^4 scattering encounters using a binary-single scattering code similar in logic to that described in Hut & Bahcall (1983). The motion was integrated with the symplectic algorithm of Mikkola & Tanikawa




FIG. 1.— Simulated outcomes for binaries with $M_1 = M_2 = 1 M_\odot$ disrupted by a $1000 M_\odot$ black hole. The panels to the top and right of the scatter plots show the histograms of $\log a_1$, $\log v_2$, and $1 - e_1$. Below the dashed curve ($a_1(1 - e_1) = 10 R_\odot$) a solar-type star is tidally disrupted by the black hole.

(1999) and Preto & Tremaine (1999). We chose a black-hole mass of $1000 M_\odot$ and stellar masses of $M_1 = M_2 = 1 M_\odot$. The binary orbits were initially circular, with separations drawn from $p(a_b) \propto a_b^{-1}$ ($a_b = 0.01$ – 1 AU). The speed at infinity was fixed at $v_\infty = 10 \text{ km s}^{-1}$. Impact parameters, b , were drawn from a distribution that is flat in $b^2 \simeq 2sG(M_\bullet + M_b)/v_\infty^2$, limited so that $s < 2r_t$. Figure 1 shows that the distributions of $\log a_1$ and $\log v_2$ are roughly flat in the middle, with high and low tails not predicted in § 2. Solar-type stars with $a_1(1 - e_1) < 10 R_\odot$ (below the dashed curve in Fig. 1) are probably tidally disrupted by the black hole (see eq. [1]).

4. DYNAMICAL EVOLUTION OF CAPTURED STARS

Each binary disruption by a $\simeq 10^3 M_\odot$ black hole deposits one star into a compact, eccentric Keplerian orbit. In 100 Myr, ~ 10 – 100 stars may be captured. However, a collection of $\gtrsim 10$ stars orbiting the black hole with periods of $\lesssim 10$ yr will not persist in the time between captures. Mechanisms that drive the evolution of captured stars include gravitational and collisional two-body encounters, and tidal disruption by the

black hole. These processes are quantified below, under the ad hoc simplifying assumptions that the stars have equal masses of $m = 1 M_\odot$, follow a power-law density profile with $\alpha \simeq 2$, and everywhere have a nearly isotropic velocity field. We neglect relatively weak and rare perturbative encounters between captured stars and cluster stars on highly elongated orbits (Gültekin, Miller, & Hamilton 2004).

Consider N point masses orbiting the black hole. Let r_h be the radius that contains half of the stars and P_h be the period of a circular orbit at r_h . The relaxation time at r_h is $\tau_h \sim 10^5 P_h M_3^2 N^{-1}$ (eq. [7] with $\Lambda = \ln(M_\bullet/m) \sim 10$). “Resonant relaxation” (Rauch & Tremaine 1996) of a star’s orbital angular momentum occurs on a shorter timescale of $\tau_h^{\text{res}} \sim 10^3 P_h M_3$, driving the collection of captured stars toward velocity isotropy without affecting the energy distribution. If $P_h \sim 10$ yr and $N \gtrsim 10$, the orbits spread significantly in the radial direction in the time between captures. Captured stars on the widest orbits will mingle with other (non-captured) stars within r_i . The outward transport of captured stars into the cusp may noticeably alter the cusp structure.

A star may be unbound from the black hole following a single strong encounter (e.g., Lin & Tremaine 1980). An upper limit to the ejection rate is obtained by setting the maximum two-body impact parameter to $b_{\text{ej}} = Gm/\sigma^2(r)$, characteristic of a $\sim 90^\circ$ deflection but not necessarily an ejection. The local rate per star is $\sim \pi n(r) b_{\text{ej}}^2 \sigma(r)$. Integrating over all captured stars, we find a rate of $\lesssim 10^{-7} P_h^{-1} M_3^{-2} N^2$. For $N \sim 10$ and $P_h \sim 10$ yr, most stars may be ejected in the time between captures. Under the point-mass assumption, energy conservation demands that ejection and relaxation leave at least one star tightly bound to the black hole. When finite stellar radii are considered, two additional processes must be addressed.

First, the (non-resonant) rate at which the black hole tidally disrupts captured stars is $\sim N/\tau_h \sim 10^{-5} P_h^{-1} M_3^{-2} N^2$, which is much more efficient than ejection at removing stars. Resonant relaxation may increase the rate substantially, since angular momentum diffusion drives stars into the loss cone. Note that if a considerable fraction (e.g., $\gtrsim 10\%$) of the captured stars are ultimately disrupted by the hole, the implied rate may exceed the swallowing rate for single stars determined by previous authors ($\sim 10^{-8} \text{ yr}^{-1}$ for $M_3 = n_5 = \sigma_{10} = 1$; e.g., Frank & Rees 1976; Lightman & Shapiro 1977). Second, the collision rate for stars of radius R is $\sim 2\pi P_h^{-1} (R/r_h)^2 N^2 \sim (10^{-9} - 10^{-6}) (R/R_\odot)^2 N^2 \text{ yr}^{-1}$ for $r_h = 10$ – 50 AU. A collision between two main-sequence stars will result in coalescence unless the relative speed at infinity is significantly larger than the escape speed from the stellar surface (e.g., Freitag & Benz 2004), which may occur within ~ 1 AU of a $10^3 M_\odot$ black hole.

When $N \gtrsim 10$ and $P_h \sim 10$ yr, all of the above processes act on timescales shorter than or comparable to the time between stellar captures. Thus, at any time, only a few stars are expected to orbit the black hole with periods of $\lesssim 10$ yr.

5. CONCLUSIONS AND OUTLOOK

A $10^3 M_\odot$ black hole at the center of a globular cluster may tidally disrupt *thousands* of binaries over 10 Gyr, each encounter leaving one star within ~ 100 AU of the hole. At any given time, the number of stars in such compact orbits is probably only a few, limited by interactions between captured stars and tidal disruption. Before we can assess possible observational consequences of these processes, numerical simulations are required to properly calculate the binary disruption rate and the dynamical evolution of captured stars.

The cluster evolution codes of Fregeau et al. (2003, Monte Carlo) and Giersz & Spurzem (2003, gas-dynamical/Monte Carlo) are well suited to the problem of determining the binary disruption rate. These codes efficiently treat a realistic number ($\gtrsim 10^5$) of stars and binaries, and include explicit integration of close binary-single and binary-binary encounters. The black hole would be represented by a central point mass, as in Freitag & Benz (2002), but the imposed spherical symmetry demands that it be fixed at the origin, neglecting its Brownian motion. These algorithms cannot adequately follow the dynamics of a small- N system of captured stars.

Direct N -body integration has been used to study the dynamics of single stars near a massive black hole (Baumgardt et al. 2004b; Preto et al. 2004). Such simulations are prohibitively time-consuming when $N \gtrsim 10^5$ and there is a sizable binary fraction. Simulations with $N \sim 10^4$ may be of value, since the dynamics of captured stars are computed correctly. It may also be useful to follow the collisional N -body evolution ~ 10 captured stars in isolation from the cluster.

The high binary disruption rate and compact orbits of captured stars increase the likelihood that a cluster IMBH will be detected, perhaps in one of the following ways:

1. If, following a stellar tidal disruption, $\simeq 0.5 M_\odot$ is accreted by the black hole with a 10% mass-to-light conversion efficiency, the duration of the Eddington-limited flare is $\sim 10^4 M_3^{-1}$ yr. Such a flare might be visible as a very bright UV/X-ray source in one of the ~ 100 globular clusters in the Milky Way if most contain a $\simeq 10^3 M_\odot$ black hole that disrupts stars at a rate of $\sim 10^{-6}$ yr $^{-1}$; but we do not, and both

suppositions may be highly optimistic.

2. If the orbit of a captured star with a pericenter distance of $\simeq 10 M_3^{1/3} R_\odot$ circularizes without the star being destroyed, stellar evolution may drive a long ($\gtrsim 10^8$ yr) phase of mass transfer, with accretion rates of $\lesssim 10^{-8} M_\odot \text{ yr}^{-1}$ and luminosities of $\lesssim 10^{38} \text{ erg s}^{-1}$ (Hopman et al. 2004; Portegies Zwart et al. 2004). Steady mass transfer from a $\gtrsim 10 M_\odot$ donor has been proposed as a fuel source for ultraluminous X-ray sources in young star clusters.

3. Hundreds of cluster neutron stars may acquire binary companions dynamically, accrete mass, and ultimately turn on as millisecond radio pulsars. A $\simeq 10^3 M_\odot$ black hole may capture 1 pulsar from a binary system every ~ 1 Gyr, making the detection of such a pulsar unlikely. The timing properties of an orbiting pulsar would clearly reveal the black hole.

4. A captured neutron star or white dwarf that migrates to within $\sim 1 R_\odot$ of a $10^3 M_\odot$ black hole would produce a detectable gravitational-wave signal for *LISA* from distances of $\lesssim 100$ Mpc in the short time ($\lesssim 100$ yr) before coalescence (see Baumgardt et al. 2004b; Portegies Zwart et al. 2004).

Special thanks go to J. Fregeau and S. Rappaport for providing valuable critical remarks on the paper. I would also like to acknowledge enlightening discussions with R. Di Stefano, S. Gaudi, A. Juett, A. Loeb, and S. Ransom. This work was supported by NASA and the Chandra Postdoctoral Fellowship program through grant number PF2-30024.

REFERENCES

- Bahcall, J. N. & Wolf, R. A. 1977, *ApJ*, 216, 883
 Barkana, R. & Loeb, A. 2001, *Phys. Rep.*, 349, 125
 Baumgardt, H., Hut, P., Makino, J., McMillan, S., & Portegies Zwart, S. 2003, *ApJ*, 582, L21
 Baumgardt, H., Makino, J., Hut, P., McMillan, S., & Portegies Zwart, S. 2003, *ApJ*, 589, L25
 Baumgardt, H., Makino, J., & Ebisuzaki, T. 2004, *ApJ*, 613, 1143
 Cohn, H. & Kulsrud, R. M. 1978, *ApJ*, 226, 1087
 Colpi, M., Possenti, A., & Gualandris, A. 2002, *ApJ*, 570, L85
 Ebisuzaki, T., et al. 2001, *ApJ*, 562, L19
 Eisenhauer, F. 2005, *ApJ*, submitted, astro-ph/0502129
 Frank, J. & Rees, M. J. 1976, *MNRAS*, 176, 633
 Fregeau, J. M., Gürkan, M. A., Joshi, K. J., & Rasio, F. A. 2003, *ApJ*, 593, 772
 Freitag, M., & Benz, W. 2002, *A&A*, 394, 345
 Freitag, M., & Benz, W. 2005, *MNRAS*, in press, astro-ph/0403621
 Gebhardt, K., Pryor, C., O'Connell, R. D., Williams, T. B., & Hesser, J. E. 2000, *AJ*, 119, 1268
 Gebhardt, K., Rich, R. M., & Ho, L. C. 2002, *ApJ*, 578, L41
 Gerssen, J., van der Marel, R. P., Gebhardt, K., Guhathakurta, P., Peterson, R. C., & Pryor, C. 2002, *AJ*, 124, 3270
 Giersz, M. & Spurzem, R. 2003, *MNRAS*, 343, 781
 Gould, A. & Quillen, A. C. 2003, *ApJ*, 592, 935
 Gültekin, K., Miller, M. C., & Hamilton, D. P. 2004, *ApJ*, submitted, astro-ph/0402532
 Hills, J. G. 1988, *Nature*, 331, 687
 Hopman, C., Portegies Zwart, S. F., & Alexander, T. 2004, *ApJ*, 604, L101
 Hut, P. & Bahcall, J. N. 1983, *ApJ*, 268, 319
 Hut, P., et al. 1992, *PASP*, 104, 981
 Islam, R. R., Taylor, J. E., & Silk, J. 2004, *MNRAS*, 365, 2966
 Ivanova, N., Belczynski, K., Fregeau, J. M., & Rasio, F. A. 2005, *MNRAS*, in press, astro-ph/0501131
 King, A. R., Davies, M. B., Ward, M. J., Fabbiano, G., & Elvis, M. 2001, *ApJ*, 552, L109
 Lightman, A. P. & Shapiro, S. L. 1977, *ApJ*, 211, 244
 Lin, D. N. C. & Tremaine, S. 1980, *ApJ*, 242, 789
 Madau, P. & Rees, M. J. 2001, *ApJ*, 551, L27
 Magorrian, J. & Tremaine, S. 1999, *MNRAS*, 309, 447
 McNamara, B. J., Harrison, T. E., & Anderson, J. 2003, *ApJ*, 595, 187
 Mikkola, S. & Tanikawa, K. 1999, *MNRAS*, 310, 745
 Miller, M. C. & Hamilton, D. P. 2002, *MNRAS*, 330, 232
 Miller, M. C. & Colbert, E. J. M. 2004, *Int. J. Mod. Phys. D*, 13, 1
 Portegies Zwart, S. F. & McMillan, S. L. W. 2002, *ApJ*, 576, 899
 Portegies Zwart, S. F., Dewi, J., & Maccarone, T. 2004, *MNRAS*, 355, 413
 Preto, M. & Tremaine, S. 1999, *AJ*, 118, 2532
 Preto, M., Merritt, D., & Spurzem, R. 2004, *ApJ*, 613, L109
 Rauch, K. P. & Tremaine, S. 1996, *New Astronomy*, 1, 149
 Rappaport, S., Podsiadlowski, Ph., & Pfahl, E. 2004, *MNRAS*, 356, 401
 Schneider, R., Ferrara, A., Natarajan, P., & Omukai, K. 2002, *ApJ*, 571, 30
 Shapiro, S. L. 1985, *IAU Symp. 113: Dynamics of Star Clusters*, 113, 373
 Spitzer, L. J. & Hart, M. H. 1971, *ApJ*, 164, 399
 Syer, D. & Ulmer, A. 1999, *MNRAS*, 306, 35
 Young, P. J. 1977, *ApJ*, 217, 287
 Yu, Q. & Tremaine, S. 2003, *ApJ*, 599, 1129